Author Index

ABO T. *see* SATO Y.
ACHA-ORBEA H. *see* PARISH N.M.
ADERKA D. *see* HERSHKOVIZ R.
AGIS H. *see* GESSL A.
AKBAR A.N. *see* GRUNDY J.E.
AKBAR S.M.F., ONJI M., INABA K., YAMAMURA K.-I. & OHTA Y. Low responsiveness of hepatitis B virus-transgenic mice in antibody response to T-cell-dependent antigen: defect in antigen-presenting activity of dendritic cells, 468
ALLÉRA A. *see* SEMMLER J.
ANELLI R., PLACIDO R., SAMBUY Y., BACH S., DI MASSIMO A. & COLIZZI V. Cytotoxic activity of human lymphocytes against differentiated intestinal tumour cell lines, 166
ARDEN B. *see* SPIEß S.
ARIZONO N. *see* YAMADA M.

BABIUK L.A. *see* DENIS M.
BACH S. *see* ANELLI R.
BADLEY R.A. *see* VERHOEYEN M.E.
BAIRD J.D. *see* JOSEPH S.
BAKER D. *see* FAIRCHILD S.
BANDO Y., KSANDER B.R. & STREILEIN J.W. Incomplete activation of lymphokine-producing T cells by alloantigenic intraocular tumours in anterior chamber-associated immune deviation, 266
BANSAL S. *see* STAINES N.A.
BARKER R.N., CASSWELL K.M. & ELSON C.J. Identification of murine erythrocyte autoantigens and cross-reactive rat antigens, 568
BARTON K., CALDER V.L. & LIGHTMAN S. Isolation of retinal lymphocytes in experimental autoimmune uveoretinitis: phenotypic and functional characterization, 393
BAZZONI F. *see* CASSATELLA M.A.
BEECHAM H.J. *see* TRAUGER R.J.
BELL E.B. *see* JAFFERY G.
BERKE G., ROSEN D. & RONEN D. Mechanism of lymphocyte-mediated cytolysis: functional cytolytic T cells lacking perforin and granzymes, 105
BETTS W.H. *see* WOZNIAK A.
BEVERLEY P.C.L. *see* WALLACE D.L.
BIENZLE U. *see* KREMSNER P.G.
BINNS R.M. *see* YANG H.
BISCHOFF P. *see* CHRIST M.
BJÖRK P. & PAULIE S. Inhibition of LFA-1-dependent human B-cell aggregation induced by CD40 antibodies and interleukin-4 leads to decreased IgE synthesis, 218
BLOKSMA N. *see* PIETERS R.H.H.
BLUE M.-L., DAVIS G., CONRAD P. & KELLEY K. Specific cleavage of the α_4 integrin associated with activation of peripheral T lymphocytes, 80
BOL M. *see* PIETERS R.H.H.
BOLTZ-NITULESCU G. *see* GESSL A.
BOSNIC M. *see* REEVE V.E.
BOUT D. *see* CHARDÈS T.
BOUVET J.-P. *see* QUAN C.P.
BRAEGGER C.P. *see* BREESE E.
BRANDON M.R. *see* GILL H.S.
BREESE E., BRAEGGER C.P., CORRIGAN C.J., WALKER-SMITH J.A. & MacDONALD T.T. Interleukin-2 and interferon-γ-secreting T cells in normal and diseased intestinal mucosa, 127
BRETT S.J. *see* YANG D.
BREW R. *see* CHRISTMAS S.E.
BRIGGS S. *see* VERHOEYEN M.E.
BRODERICK E.L. *see* VERHOEYEN M.E.
BRUNNER H. *see* SCHLÜTER D.
BURNETT K.G. *see* TRAUGER R.J.
BUTCHER G.W. *see* JOSEPH S.
BUZONI-GATEL D. *see* CHARDÈS T.

CALDER V.L. *see* BARTON K.
CALLANAN J.J. *see* WILLETT B.J.
CALLOW M.G. *see* HOYNE G.F.
CAMBIASO C.L. *see* LUCAS G.P.
CARLO D.J. *see* TRAUGER R.J.
CARTER R. *see* NAOTUNNE T. DE S.
CASSATELLA M.A., GUASPARRI I., CESKA M., BAZZONI F. & ROSSI F. Interferon-gamma inhibits interleukin-8 production by human polymorphonuclear leucocytes, 177
CASSWELL K.M. *see* BARKER R.N.
CESKA M. *see* CASSATELLA M.A.
CEUPPENS J.L. *see* VANDENBERGHE P.
CHADEE K. *see* LIN J.-Y.
CHARDÈS T., VELGE-ROUSSEL F., MEVELEC P., MEVELEC M.-N., BUZONI-GATEL D. & BOUT D. Mucosal and systematic cellular immune responses induced by *Toxoplasma gondii* antigens in cyst orally infected mice, 421
CHARTERIS D.G. & LIGHTMAN S.L. *In vivo* lymphokine production in experimental autoimmune uveoretinitis, 387
CHAVES M.F. *see* KREMSNER P.G.
CHEN Y. *see* XIANG J.
CHRIST M., LUU B., MEJIA J.E., MOOSBRUGGER I. & BISCHOFF P. Apoptosis induced by oxysterols in murine lymphoma cells and in normal thymocytes, 455
CHRISTMAS S.E., BREW R., DENIZ G. & TAYLOR J.J. T-cell receptor heterogeneity of $\gamma\delta$ T-cell clones from human female reproductive tissues, 436
CLAESSON M.H., DISSING S., TSCHERNING T. & GEISLER C. T-cell activation. V. Anti-major histocompatibility complex class I antibody-induced activation and clonal abortion in Jurkat T-leukaemic cells, 444
CLARK-LEWIS I. *see* SMITH W.B.
COHEN I.R. *see* HERSHKOVIZ R.
COLEMAN J.W. *see* HOLLIDAY M.R.
see also JAFFERY G.
COLIZZI V. *see* ANELLI R.
CONRAD P. *see* BLUE M.-L.
COOKE A. *see* PARISH N.M.
CORRIGAN C.J. *see* BREESE E.
CUNHA F.Q., MOSS D.W., LEAL L.M.C.C., MONCADA S. & LIEW F.Y. Induction of macrophage parasiticidal activity by *Staphylococcus aureus* and exotoxins through the nitric oxide synthesis pathway, 563

DAVIS G. *see* BLUE M.-L.
DE GIORGI L. *see* MATOSSIAN-ROGERS A.
DEARMAN R.J. *see* HOLLIDAY M.R.
DECKERT-SCHLÜTER M. *see* SCHLÜTER D.
DELLEMIJN T.A.M. *see* VYTH-DREESE F.A.
DENBURY A.N. *see* STAINES N.A.
DENIS M., SLAOUI M., KEIL G., BABIUK L.A., ERNST E., PASTORET P.-P. & THIRY E. Identification of different target glycoproteins for bovine herpes virus type 1-specific cytotoxic T lymphocytes depending on the method of *in vitro* stimulation, 7
DENIZ G. *see* CHRISTMAS S.E.
DI MASSIMO A. *see* ANELLI R.
DIAMOND A.G. *see* JOSEPH S.
DIAZ-SANCHEZ D., LEE T.H. & KEMENY D.M. Ricin enhances IgE responses by inhibiting a subpopulation of early-activated IgE regulatory $CD8^+$ T cells, 226
DIAZ-SANCHEZ D., NOBLE A., STAYNOV D.Z., LEE T.H. & KEMENY D.M. Elimination of IgE regulatory rat $CD8^+$ T cells *in vivo* differentially modulates interleukin-4 and interferon-γ but not interleukin-2 production by splenic T cells, 513
DIOCEE M. *see* RAYNER D.C.
DISSING S. *see* CLAESSON M.H.
DOE W.F. *see* PAVLI P.
DOWNES K.L. *see* GRUNDY J.E.

DRAKE W.T. & ISSEKUTZ A.C. Transforming growth factor-β_1 enhances polymorphonuclear leucocyte accumulation in dermal inflammation and transendothelial migration by a priming action, 197
DUFFY P.C. *see* TRAUGER R.J.

EIDA S.J. *see* VERHOEYEN M.E.
EILAT D. *see* STAINES N.A.
EISENHUT T. *see* SEMMLER J.
ELSON C.J. *see* BARKER R.N.
ENDRES S. *see* SEMMLER J.
ENNEN J. & KURTH R. Interferon-γ-activated monocytes impair infectivity of HIV particles by an oxygen metabolite-dependent reaction, 171
ERNST E. *see* DENIS M.
ETO M. *see* MAEDA T.

FAIRCHILD S., BAKER D. & TURK J.L. Characterization of experimental allergic encephalomyelitis-susceptible Biozzi AB/H (H-2^{dq1}) mice which express H-$2A^{nod}$: analysis of T-cell receptor expression and the detection of a deletion ligand encoded by Mtv-7, 260
FALUS A. Interleukin-6 biosynthesis is increased by histamine in human B-cell and glioblastoma cell lines, 193
FARIA A.M.C., GARCIA G., RIOS M.J.C., MICHALAROS C.L. & VAZ N.M. Decrease in susceptibility to oral tolerance induction and occurrence of oral immunization to ovalbumin in 20–38-week-old mice. The effect of interval between oral exposures and rate of antigen intake in the oral immunization, 147
FERRE F. *see* TRAUGER R.J.
FIGDOR C.G. *see* VYTH-DREESE F.A.
FILSINGER S. *see* JAHN B.
FLORA P.S. *see* RAPLEY R.
FÖRSTER O. *see* GESSL A.
FUJISAWA Y. *see* HAZAMA M.
FUKUHARA Y. *see* TAKENAKA T.

GAMBLE J.R. *see* SMITH W.B.
GARCIA G. *see* FARIA A.M.C.
GARSIDE P. & MOWAT A.MCI. Natural killer cells and tumour necrosis factor-α-mediated enteropathy in mice, 335
GEBERT U. *see* SEMMLER J.
GEISLER C. *see* CLAESSON M.H.
GESSL A., WILLHEIM M., AGIS H., SPITTLER A., SCHEDLE A., KRUGLUGER W., FÖRSTER O. & BOLTZ-NITULESCU G. Tumour necrosis factor-α augments the expression of Fc IgE receptor (FcεRII/CD23) on human monocytic cell lines and down-regulates interleukin-4-driven FcεRII expression on monocytes, 476
GIBBONS A.E. *see* PRICE P.
GIERMAKOWSKA W.K. *see* TRAUGER R.J.
GILAT D. *see* HERSHKOVIZ R.
GILL H.S., WATSON D.L. & BRANDON M.R. Monoclonal antibody to $CD4^+$ T cells abrogates genetic resistance to *Haemonchus contortus* in sheep, 43
GOODACRE J.A., MIDDLETON S., LYNN S., ROSS D.A. & PEARSON J. Human cartilage aggrecan CS1 region contains cryptic T-cell recognition sites, 586
GRADEHANDT G., KLEBER N., MATTNER F., MILBRADT S. & RÜDE E. The I-A^b-restricted alloresponse of D10.G4.1 T cells is based on the recognition of an endogenous peptide, 592
GRAU G.E. *see* KREMSNER P.G.
GRUNDY J.E. & DOWNES K.L. Up-regulation of LFA-3 and ICAM-1 on the surface of fibroblasts infected with cytomegalovirus, 405
GRUNDY J.E., PAHAL G.S. & AKBAR A.N. Increased adherence of CD2 peripheral blood lymphocytes to cytomegalovirus-infected fibroblasts is blocked by anti-LFA-3 antibody, 413
GUASPARRI I. *see* CASSATELLA M.A.

HÄNSCH G.M. *see* JAHN B.
HARADA T. *see* TORII I.
HARRIS T. *see* STEWART A.G.
HARRISON R.A. *see* VAN DEN BERG C.W.
HARTLEY O. *see* STAINES N.A.
HAYGLASS K.T. *see* YANG X.
HAZAMA M., MAYUMI-ANONO A., MIYAZAKI T., HINUMA S. & FUJISAWA Y. Intranasal immunization against herpes simplex virus infection by using a recombinant glycoprotein D fused with immunomodulating proteins, the B subunit of *Escherichia coli* heat-labile enterotoxin and interleukin-2, 643
HERSHKOVIZ R., GILAT D., MIRON S., MEKORI Y.A., ADERKA D., WALLACH D., VLODAVSKY I., COHEN I.R. & LIDER O. Extracellular matrix induces tumour necrosis factor-α secretion by an interaction between resting $CD4^+$ T cells and macrophages, 50
HINUMA S. *see* HAZAMA M.
HIROMATSU K. *see* MATSUZAKI G.
HOF H. *see* SCHLÜTER D.
HOGARTH P.M. *see* QUILLIAM A.L.
HOLLIDAY M.R., DEARMAN R.J., KIMBER I. & COLEMAN J.W. Sensitization of mice to chemical allergens modulates the responsiveness of isolated mast cells to IgE-dependent activation, 508
HORII Y. *see* ISHIKAWA N.
HOSIE M.J. *see* WILLETT B.J.
HOYNE G.F., CALLOW M.G., KUHLMAN J. & THOMAS W.R. T-cell lymphokine response to orally administered proteins during priming and unresponsiveness, 534
HOYNE G.F., CALLOW M.G., KUO M.-C. & THOMAS W.R. Characterization of T-cell responses to the house dust mite allergen *Der p* II in mice. Evidence for major and cryptic epitopes, 65
HOYNE G.F., CALLOW M.G., KUO M.-C. & THOMAS W.R. Comparison of antigen presentation by lymph node cells from protein and peptide-primed mice, 58
HUME D.A. *see* PAVLI P.
HVIID L., ØDUM N. & THEANDER T.G. The relation between T-cell expression of LFA-1 and immunological memory, 237

IIAI T. *see* SATO Y.
IMAEDA H. *see* MIURA S.
INABA K. *see* AKBAR S.M.F.
ISENBERG D.A. *see* STAINES N.A.
ISHIKAWA N., HORII Y. & NAWA Y. Immune-mediated alteration of the terminal sugars of goblet cell mucins in the small intestine of *Nippostrongylus brasiliensis*-infected rats, 303
ISSEKUTZ A.C. *see* DRAKE W.T.

JACKSON J. *see* MALHOTRA R.
JAFFERY G., BELL E.B. & COLEMAN J.W. Induction of an auto-anti-IgE response in rats. IV. Effects on mast cell degranulation, 635
JAHN B., VON KEMPIS J., KRÄMER K.-L., FILSINGER S. & HÄNSCH G.M. Interaction of the terminal complement components C5b–9 with synovial fibroblasts: binding to the membrane surface leads to increased levels in collagenase-specific mRNA, 329
JARRETT O. *see* WILLETT B.J.
JENSEN F.C. *see* TRAUGER R.J.
JENSENIUS J.-C. *see* MALHOTRA R.
JOSEPH S., DIAMOND A.G., SMITH W., BAIRD J.D. & BUTCHER G.W. BB-DR/Edinburgh: a lymphopenic, non-diabetic subline of BB rats, 318

KAHAN M. *see* QUARATINO S.
KARUNAWEERA *see* NAOTUNNE T. DE S.
KEES U.R. *see* SALVATI P.D.
KEIL G. *see* DENIS M.
KELLER K. *see* LIN J.-Y.
KELLEY K. *see* BLUE M.-L.
KELSO A. *see* THOMSON J.A.
KEMENY D. M. *see* DIAZ-SANCHEZ D.
KIMBER I. *see* HOLLIDAY M.R.
KITAMURA Y. *see* TORII I.
KLEBER N. *see* GRADEHANDT G.
KODAMA T. *see* SAGAWA T.
KONG Y. -Y. *see* MAEDA T.
KÖNIG B. & KÖNIG W. Induction and suppression of cytokine release (tumour necrosis factor-α; interleukin-6, interleukin-1β) by *Escherichia coli* pathogenicity factors (adhesions, α-haemolysin), 526

KÖNIG W. *see* KÖNIG B.
KOROPATNICK J. *see* XIANG J.
KRÄMER K.-L. *see* JAHN B.
KREMSNER P.G., NÜSSLER A., NEIFER S., CHAVES M.F., BIENZLE U., SENALDI G. & GRAU G.E. Malaria antigen and cytokine-induced production of reactive nitrogen intermediates by murine macrophages: no relevance to the development of experimental cerebral malaria, 286
KROEGEL C. & MATTHYS H. Platelet-activating factor-induced human eosinophil activation. Generation and release of cyclooxygenase metabolites in human blood eosinophils from asthmatics, 279
KRUGLUGER W. *see* GESSL A.
KSANDER B.R. *see* BANDO Y.
KUHLMAN J. *see* HOYNE G.F.
KUHRÖBER A. *see* SPIEß S.
KUO M.-C. *see* HOYNE G.F.
KURIBAYASHI K. *see* TAKENAKA T.
KURTH R. *see* ENNEN J.

LAM B.W. *see* PIETERS R.H.H.
LEAL L.M.C.C. *see* CUNHA F.Q.
LEE T.H. *see* DIAZ-SANCHEZ D.
LEWIS D.E. *see* TRAUGER R.J.
LI K. *see* XIANG J.
LI X.-Y., MATSUZAKI G., YOSHIKAI Y., MURAMORI K. & NOMOTO K. T cells expressing both L-selectin and CD44 molecules increase in number in peritoneal exudate cells and *in vitro*-stimulated spleen cells from mice immunized intraperitoneally with *Listeria monocytogenes*, 28
LIDER O. *see* HERSHKOVIZ R.
LIEW F.Y. *see* CUNHA F.Q.
see also YANG D.
LIGHTMAN S. *see* BARTON K.
LIGHTMAN S.L. *see* CHARTERIS D.G.
LIM B.L., LU J. & REID K.B.M. Structural similarity between bovine conglutinin and bovine lung surfactant protein D and demonstration of liver as a site of synthesis of conglutinin, 159
LIN J.-Y., KELLER K. & CHADEE K. *Entamoeba histolytica* proteins modulate the respiratory burst potential by murine macrophages, 291
LONDEI M. *see* QUARATINO S.
LOWRIE D.B. *see* SILVA C.L.
LU J. *see* LIM B.L.
LUCAS G.P., CAMBIASO C.L. & VAERMAN J.P. An interspecies idiotope associated with the anti-cholera toxin response detected by a monoclonal auto-anti-idiotypic antibody, 498
LUKACS K. *see* SILVA C.L.
LUND T. *see* PARISH N.M.
LUO X. *see* XIANG J.
LUU B. *see* CHRIST M.
LYNN S. *see* GOODACRE J.A.

MacDONALD T.T. *see* BREESE E.
McKENZIE I.F.C. *see* QUILLIAM A.L.
McLENNAN G. *see* WOZNIAK A.
MAEDA J. *see* TAKENAKA T.
MAEDA T., ETO M., NISHIMURA Y., NOMOTO K., KONG Y.-Y. & NOMOTO K. Direct evidence for clonal destruction of alloreactive T cells in the mice treated with cyclophosphamide after allopriming, 113
MAHLMAN R.T. *see* PAPPO J.
MALHOTRA R., WILLIS A.C., JENSENIUS J.-C., JACKSON J. & SIM R.B. Structure and homology of human C1q receptor (collectin receptor), 341
MARUGG J.D. *see* VERHOEYEN M.E.
MATOSSIAN-ROGERS A. & DE GIORGI L. Anti-T-cell receptor Vβ6 breaks tolerance in Mls^a mice and induces production of anti-Mls^a antibodies, 122
MATSUMOTO Y. *see* SATO Y.
MATSUO T. *see* QUAN C.P.
MATSUZAKI G. *see* LI X.-Y.
MATSUZAKI G., HIROMATSU K., YOSHIKAI Y., MURAMORI K. & NOMOTO K. Characterization of T-cell receptor $\gamma\delta$ T cells appearing at the early phase of murine *Listeria monocytogenes* infection, 22
MATTHYS H. *see* KROEGEL C.
MATTNER F. *see* GRADEHANDT G.
MAYUMI-ANONO A. *see* HAZAMA M.
MEJIA J.E. *see* CHRIST M.
MEKORI Y.A. *see* HERSHKOVIZ R.
MELIEF C.J.M. *see* VYTH-DREESE F.A.
MENDIS K.N. *see* NAOTUNNE T.DE S.
MEVELEC M.-N. *see* CHARDÈS T.
MEVELEC P. *see* CHARDÈS T.
MICHALAROS C.L. *see* FARIA A.M.C.
MIDDLETON S. *see* GOODACRE J.A.
MIHARA M. *see* TAKENAKA T.
MILBRADT S. *see* GRADEHANDT G.
MIRON S. *see* HERSHKOVIZ R.
MITCHINER J. *see* STAINES N.A.
MIURA S., IMAEDA H., SHIOZAKI H., OHKUBO N., TASHIRO H., SERIZAWA H., TSUCHIYA M. & TSO P. Increased proliferative response of lymphocytes from intestinal lymph during long chain fatty acid absorption, 142
MIYAZAKI T. *see* HAZAMA M.
MOELLER J. *see* SEMMLER J.
MONCADA S. *see* CUNHA F.Q.
MOOREN A.T.A. *see* VERHOEYEN M.E.
MOOSBRUGGER I. *see* CHRIST M.
MORGAN B.P. *see* VAN DEN BERG C.W.
MORIKAWA S. *see* TORII I.
MOSS D.W. *see* CUNHA F.Q.
MOWAT A.MCI. *see* GARSIDE P.
MOYANA T. *see* XIANG J.
MURAMORI K. *see* LI X.-Y.
see also MATSUZAKI G.
MUTO T. *see* SATO Y.

NAKAMINE H. *see* TAKENAKA T.
NAKAZAWA M. *see* YAMADA M.
NAOTUNNE T. DE S., KARUNAWEERA, MENDIS K.N. & CARTER R. Cytokine-mediated inactivation of malarial gametocytes is dependent on the presence of white blood cells and involves reactive nitrogen intermediates, 555
NAWA Y. *see* ISHIKAWA N.
NEIFER S. *see* KREMSNER P.G.
NEIL J.C. *see* WILLETT B.J.
NEWMAN I. & WILKINSON P.C. Locomotor responses of human CD45 lymphocyte subsets: preferential locomotion of CD45RO$^+$ lymphocytes in response to attractants and mitogens, 92
NISHIMURA Y. *see* MAEDA T.
NOBLE A. *see* DIAZ-SANCHEZ D.
NOMOTO K. *see* LI X.-Y.
see also MAEDA T.
see also MATSUZAKI G.
NÜSSLER A. *see* KREMSNER P.G.

ØDUM N. *see* HVIID L.
OHKUBO N. *see* MIURA S.
OHMORI K. *see* SATO Y.
OHSUGI Y. *see* TAKENAKA T.
OHTA Y. *see* AKBAR S.M.F.
OKADA M. *see* SAGAWA T.
OLVER S.D. *see* PRICE P.
ONJI M. *see* AKBAR S.M.F.
OSMAN N. *see* QUILLIAM A.L.

PAHAL G.S. *see* GRUNDY J.E.
PAPPO J. & MAHLMAN R.T. Follicle epithelial M cells are a source of interleukin-1 in Peyer's patches, 505
PARISH N.M., ACHA-ORBEA H., SIMPSON E., QIN S.-X., LUND T. & COOKE A. A comparative study of T-cell receptor Vβ usage in non-obese diabetic (NOD) and I-E transgenic NOD mice, 606
PASTORET P.-P. *see* DENIS M.
PAULIE S. *see* BJÖRK P.
PAVLI P., HUME D.A., VAN DE POL E. & DOE W.F. Dendritic cells, the major antigen-presenting cells of the human colonic lamina propria, 132

PEARSON J. *see* GOODACRE J.A.
PENNINKS A.H. *see* PIETERS R.H.H.
PERONI S.E. *see* SALVATI P.D.
PETRYCKY-COX L.D. *see* RAYNER D.C.
PIETERS R.H.H., BOL M., LAM B.W., SEINEN W., BLOKSMA N. & PENNINKS A.H. Recovery from chemically induced thymus atrophy starts with $CD4^-$ $CD8^-$ $CD2^{high}$ $TcR\alpha\beta^{-/low}$ thymocytes and results in an increased formation of $CD4^-$ $CD8^-$ $TcR\alpha\beta^{high}$ thymocytes, 616
PILLOT J. *see* QUAN C.P.
PIRES R. *see* QUAN C.P.
PLACIDO R. *see* ANELLI R.
PRICE M.R. *see* VERHOEYEN M.E.
PRICE P., OLVER S.D., GIBBONS A.E. & SHELLAM G.R. B-cell activation following murine cytomegalovirus infection: implications for autoimmunity, 14

QI Y. *see* XIANG J.
QIN S.-X. *see* PARISH N.M.
QUAN C.P., WATANABE S., VUILLIER F., PIRES R., MATSUO T., STANISLAWSKI M., PILLOT J. & BOUVET J.-P. Purification and partial amino acid sequence of suppressive lymphokine from a $CD8^+$ $CD57^+$ human T hybridoma, 205
QUARATINO S., VERHOEF A., KAHAN M. & LONDEI M. Expression of the identical $V\beta$ gene in human T-cell clones does not confer the same pattern of responsiveness to bacterial enterotoxins, 600
QUILLIAM A.L., OSMAN N., McKENZIE I.F.C. & HOGARTH P.M. Biochemical characterization of murine $Fc\gamma RI$, 358

RANFORD P.R. *see* SALVATI P.D.
RAPLEY R., FLORA P.S., WALSH D.J. & WALKER M.R. Nucleotide sequence analysis of CDR3 elements of a panel of anti-peptide monoclonal antibodies recognizing parathyroid hormone-related protein, 379
RAYNER D.C., PETRYCKY-COX L.D., DIOCEE M. & RECTOR E. Major histocompatibility complex class II-restricted cytotoxicity by self-myelin basic protein-reactive T-cell hybridomas: evidence for a tumour necrosis factor-independent nucleolytic mechanism, 273
RECTOR E. *see* RAYNER D.C.
REEVE V.E., BOSNIC M. & ROZINOVA E. Carnosine (β-alanylhistidine) protects from the suppression of contact hypersensitivity by ultraviolet B (280–320 nm) radiation or by *cis* urocanic acid, 99
REID K.B.M. *see* LIM B.L.
REIMANN J. *see* SPIEß S.
RIOS M.J.C. *see* FARIA A.M.C.
ROGERS M.V. *see* YANG D.
RONEN D. *see* BERKE G.
ROSEN D. *see* BERKE G.
ROSS D.A. *see* GOODACRE J.A.
ROSSI F. *see* CASSATELLA M.A.
ROZINOVA E. *see* REEVE V.E.
RUBIO N. *see* SIERRA A.
RÜDE E. *see* GRADEHANDT G.

SAGAWA T., TOMINAGA A., KODAMA T. & OKADA M. Cytotoxicity of unstimulated and thrombin-activated platelets to human tumour cells, 650
SALVATI P.D., PERONI S.E., RANFORD P.R. & KEES U.R. Requirements for the induction of the interleukin-2 receptor complex in a human pre-T-cell line, 449
SAMBUY Y. *see* ANELLI R.
SATO Y., TSUKADA K., IIAI T., OHMORI K., YOSHIDA K., MUTO T., WATANABE H., MATSUMOTO Y. & ABO T. Activation of extrathymic T cells in the liver during liver regeneration following partial hepatectomy, 86
SAUNDERS J.A. *see* VERHOEYEN M.E.
SCHEDLE A. *see* GESSL A.
SCHIRMBECK R. *see* SPIEß S.
SCHLÜTER D., DECKERT-SCHLÜTER M., SCHWENDEMANN G., BRUNNER H. & HOF H. Expression of major histocompatibility complex class II antigens and levels of interferon-γ, tumour necrosis factor, and interleukin-6 in cerebrospinal fluid and serum in *Toxoplasma gondii*-infected SCID and immunocompetent C.B-17 mice, 430
SCHÖNHARTING M.M. *see* SEMMLER J.
SCHWENDEMANN G. *see* SCHLÜTER D.
SCICCHITANO R. *see* WOZNIAK A.
SEINEN W. *see* PIETERS R.H.H.
SEMMLER J., GEBERT U., EISENHUT T., MOELLER J., SCHÖNHARTING M.M., ALLÉRAA. & ENDRES S. Xanthine derivatives: comparison between suppression of tumour necrosis factor-α production and inhibition of cAMP phosphodiesterase activity, 520
SENALDI G. *see* KREMSNER P.G.
SERIZAWA H. *see* MIURA S.
SHELLAM G.R. *see* PRICE P.
SHIOZAKI H. *see* MIURA S.
SIERRA A. & RUBIO N. Theiler's murine encephalomyelitis virus induces tumour necrosis factor-α in murine astrocyte cell cultures, 399
SILVA C.L., LUKACS K. & LOWRIE D.B. Major histocompatibility complex non-restricted presentation to $CD4^+$ T lymphocytes of *Mycobacterium leprae* heat-shock protein 65 antigen by macrophages transfected with the mycobacterial gene, 35
SIM R.B. *see* MALHOTRA R.
SIMPSON E. *see* PARISH N.M.
SLAOUI M. *see* DENIS M.
SMITH W. *see* JOSEPH S.
SMITH W.B., GAMBLE J.R., CLARK-LEWIS I. & VADAS M.A. Chemotactic desensitization of neutrophils demonstrates interleukin-8 (IL-8)-dependent and IL-8-independent mechanisms of transmigration through cytokine-activated endothelium, 491
SOO C. *see* SULLIVAN D.A.
SPIEß S., KUHRÖBER A., SCHIRMBECK R., ARDEN B. & REIMANN J. Diversity of functional T-cell receptor δ-chain transcripts from bone marrow cells of athymic nude mice, 252
SPITTLER A. *see* GESSL A.
STAINES N.A., WARD F.J., DENBURY A.N., MITCHINER J., HARTLEY O., EILAT D., ISENBERG D.A. & BANSAL S. Primary sequence and location of V-88, a DNA-binding monoclonal autoantibody, determined by idiotope scanning with synthetic peptides on pins, 371
STANISLAWSKI M. *see* QUAN C.P.
STAYNOV D.Z. *see* DIAZ-SANCHEZ D.
STEINIGER B. & VAN DER MEIDE P.H. High-dose interferon-γ alters the distribution of B lymphocytes and macrophages in rat spleen and lymph nodes, 461
STEWART A.G. & HARRIS T. Adenosine inhibits platelet-activating factor, but not tumour necrosis factor-α-induced priming of human neutrophils, 152
STREILEIN J.W. *see* BANDO Y.
SULLIVAN D.A., VAERMAN J.-P. & SOO C. Influence of severe protein malnutrition on rat lacrimal, salivary and gastrointestinal immune expression during development, adulthood and ageing, 308

TAKENAKA T., KURIBAYASHI K., NAKAMINE H., TSUJIMOTO M., FUKUHARA Y., MAEDA J., MIHARA M., UCHIYAMA Y. & OHSUGI Y. Regulation by cytokines of eosinophilopoiesis and immunoglobulin E production in mice, 541
TASHIRO H. *see* MIURA S.
TAYLOR J.J. *see* CHRISTMAS S.E.
THEANDER T.G. *see* HVIID L.
THIRY E. *see* DENIS M.
THOMAS W.R. *see* HOYNE G.F.
THOMSON J.A., TROUTT A.B. & KELSO A. Contact sensitization to oxazolone: involvement of both interferon-γ and interleukin-4 in oxazolone-specific Ig and T-cell responses, 185
TOMINAGA A. *see* SAGAWA T.
TORII I., MORIKAWA S., HARADA T. & KITAMURA Y. Two distinct types of cellular mechanisms in the development of delayed hypersensitivity in mice: requirement of either mast cells or macrophages for elicitation of the response, 482
TRAUGER R.J., GIERMAKOWSKA W.K., FERRE F., DUFFY P.C., WALLACE M.R., LEWIS D.E., BEECHAM H.J., BURNETT K.G., JENSEN F.C. & CARLO D.J. Cell-mediated immunity of HIV-1 in Walter Reed stages 1–6 individuals: correlation with virus burden, 611
TROUTT A.B. *see* THOMSON J.A.
TSCHERNING T. *see* CLAESSON M.H.

TSO P. *see* MIURA S.
TSUCHIYA M. *see* MIURA S.
TSUJIMOTO M. *see* TAKENAKA T.
TSUKADA K. *see* SATO Y.
TURK J.L. *see* FAIRCHILD S.

UCHIYAMA Y. *see* TAKENAKA T.

VADAS M.A. *see* SMITH W.B.
VAERMAN J.-P. *see* SULLIVAN D.A.
VAERMAN J.P. *see* LUCAS G.P.
VAN DE POL E. *see* PAVLI P.
VAN DEN BERG C.W., HARRISON R.A. & MORGAN B.P. The sheep analogue of human CD59: purification and characterization of its complement inhibitory activity, 349
VAN DER MEIDE P.H. *see* STEINIGER B.
VAN KOOYK Y. *see* VYTH-DREESE F.A.
VAN VAECK F. *see* VANDENBERGHE P.
VANDENBERGHE P., VERWILGHEN J., VAN VAECK F. & CEUPPENS J.L. Ligation of the CD5 or CD28 molecules on resting human T cells induces expression of the early activation antigen CD69 by a calcium- and tyrosine kinase-dependent mechanism, 210
VAZ N.M. *see* FARIA A.M.C.
VELGE-ROUSSEL F. *see* CHARDÈS T.
VERHOEF A. *see* QUARATINO S.
VERHOEYEN M.E., SAUNDERS J.A., PRICE M.R., MARUGG J.D., BRIGGS S., BRODERICK E.L., EIDA S.J., MOOREN A.T.A. & BADLEY R.A. Construction of a reshaped HMFG1 antibody and comparison of its fine specificty with that of the parent mouse antibody, 364
VERWILGHEN J. *see* VANDENBERGHE P.
VLODAVSKY I. *see* HERSHKOVIZ R.
VON KEMPIS J. *see* JAHN B.
VUILLIER F. *see* QUAN C.P.
VYTH-DREESE F.A., VAN KOOYK Y., DELLEMIJN T.A.M., MELIEF C.J.M. & FIGDOR C.G. Regulatory effect of interleukin-4 (IL-4) on the expression and function of lymphocyte adhesion receptors involved in IL-2-induced cell aggregation, 244

WALKER M.R. *see* RAPLEY R.
WALKER-SMITH J.A. *see* BREESE E.
WALLACE D.L. & BEVERLEY P.C.L. Characterization of a novel subset of T cells from human spleen that lacks L-selectin, 623
WALLACE M.R. *see* TRAUGER R.J.
WALLACH D. *see* HERSHKOVIZ R.
WALSH D.J. *see* RAPLEY R.
WARD F.J. *see* STAINES N.A.
WATANABE H. *see* SATO Y.
WATANABE S. *see* QUAN C.P.
WATSON D.L. *see* GILL H.S.
WILKINSON P.C. *see* NEWMAN I.
WILLETT B.J., HOSIE M.J., CALLANAN J.J., NEIL J.C. & JARRETT O. Infection with feline immunodeficiency virus is followed by the rapid expansion of a $CD8^+$ lymphocyte subset, 1
WILLHEIM M. *see* GESSL A.
WILLIS A.C. *see* MALHOTRA R.
WOZNIAK A., BETTS W.H., McLENNAN G. & SCICCHITANO R. Activation of human neutrophils by tachykinins: effect on formyl-methionyl-leucyl-phenylalaline- and platelet-activating factor-stimulated superoxide anion production and antibody-dependent cell-mediated cytotoxicity, 629

XIANG J., KOROPATNICK J., QI Y., LUO X., MOYANA T., LI K. & CHEN Y. Production of a bifunctional hybrid molecule B72.3/metallothionein-1 by protein engineering, 574

YAMADA M., NAKAZAWA M. & ARIZONO N. IgE and IgG2a antibody responses are induced by different antigen groups of the nematode *Nippostrongylus brasiliensis* in rats, 298
YAMAMURA K.-I. *see* AKBAR S.M.F.
YANG D., ROGERS M.V., BRETT S.J. & LIEW F.Y. Immunological analysis of the zinc-binding peptides of surface metalloproteinase (gp63) of *Leishmania major*, 582
YANG H. & BINNS R.M. Isolation and characterization of the soluble and membrane-bound porcine CD44 molecules, 547
YANG X. & HAYGLASS K.T. Allergen-dependent induction of interleukin-4 synthesis *in vivo*, 74
YOSHIDA K. *see* SATO Y.
YOSHIKAI Y. *see* LI X.-Y.
see also MATSUZAKI G.

Subject Index

Adenosine:
inhibition of platelet-activating factor but not tumour necrosis factor-α priming of neutrophils, 152
Alloresponse, I-A^b restricted:
based on recognition of endogenous peptide, 592
Antibody responses to antigens of *Nippostrongylus brasiliensis*, 298
Antigen presentation:
by lymph node cells from protein- and peptide-primed mice, 58
Apoptosis:
induced by oxysterols, 455
Auto-anti-IgE response:
effects on mast cell degranulation, 635
Autoimmune uveoretinitis, experimental:
phenotypic and functional characterization of retinal lymphocytes, 393

B-cell activation:
in murine cytomegalovirus infection, 14
B-cell aggregation, LFA-1 dependent:
inhibition by anti-CD40 and interleukin-4, 218
BB-DR/Edinburgh:
lymphopenic non-diabetic subline of BB rats, 318

C1q receptor, human:
structure of, 341
Cartilage aggrecan, human:
T-cell sites, 586
$CD2^+$ lymphocytes:
adherence to cytomegalovirus-infected cells, 413
CD44, porcine:
characterization of, 547
CD59, sheep analogue of human:
purification and characterization, 349
CDR3:
analysis of panel of anti-PTHrP monoclonal antibodies, 379
Complement C5b–9:
interaction with synovial fibroblasts, 329
Conglutinin, bovine:
structural similarity with bovine lung surfactant protein D, 159
Contact hypersensitivity suppression, by UVB radiation, or *cis* urocanic acid:
protection by carnosine, 99
Contact sensitization, to oxazolone:
role of interferon-γ and interleukin-4, 185
Cyclophosphamide-induced clonal destruction, 113
Cytokine release:
by *Escherichia coli* pathogenicity factors, 526
Cytokine-mediated gametocyte inactivation, 555
Cytokines:
regulation of eosinophilopoiesis and IgE production, 541
Cytomegalovirus:
increases cell surface LFA-3 and ICAM-1, 405
Cytotoxic T cells, to bovine herpes virus type 1:
target glycoproteins, 7
Cytotoxicity, major histocompatibility complex class-restricted:
evidence for tumour necrosis factor-independent nucleolytic mechanism, 273

Dendritic cells, defective:
in hepatitis B transgenic mice, 468
Dendritic cells, human colonic:
antigen presenting, 132
Delayed hypersensitivity:
requirement for mast cells or macrophages, 482

Enterotoxins, bacterial:
T-cell responsiveness, 600
Erythrocyte autoantigen:
murine, 568
Extrathymic T cells:
and liver regeneration, 86

FcγRI:
biochemical characterization, 358
Feline immunodeficiency virus infection:
expansion of $CD8^+$ lymphocytes, 1

Gamma-delta (γδ) T-cell clones:
from human female reproductive tissues, 436
Gamma-delta (γδ) T cells, in listeriosis:
V gene usage, 22

Haemonchosis, genetic resistance to:
abrogation by anti-CD4, 43
HIV infectivity:
inactivation by interferon-γ-activated monocytes, 171
HIV-1, cell-mediated immunity to:
correlation with virus burden, 611
Hybrid antibody:
metallothionein molecule, 574

Idiotope, interspecies, associated with anti-cholera toxin:
detected by auto-anti-idiotypic antibody, 498
Idiotype mapping:
of a DNA-binding monoclonal autoantibody, 371
IgE regulatory $CD8^+$ T cells, rat:
modulation of cytokine production, 513
Integrin α_4:
proteolytic cleavage on T cells, 80
Interferon-γ:
effects on distribution of B lymphocytes and macrophages, 461
inhibition of interleukin-8 production by human polymorphonuclear leucocytes, 177
Interleukin-2 receptor:
induction of, 449
Interleukin-4:
regulatory effect of interleukin-2-induced cell aggregation, 244
Interleukin-4, *in vivo* synthesis:
allergen-dependent induction, 74
Interleukin-6 biosynthesis, in B cells and glioblastoma cells: increased by histamine, 193
Intranasal immunization:
against herpes simplex virus, 643

L-selectin $CD44^+$ T cells:
after listeria immunization, 28
L-selectin negative T-cell subsets in human spleen, 629
Ligation of CD5 and CD28:
induction of CD69 expression, 210
Locomotor responses of human CD45 lymphocyte subsets, 92
Lymphocyte function-associated antigen-1 expression, on T cells:
and memory, 237
Lymphocyte proliferation:
during fatty acid absorption, 142
Lymphokines:
in gut disease, 127

M cells:
source of interleukin-1, 505
Macrophage parasiticidal activity, induction:
through the nitric oxide pathway, 563
Major histocompatibility complex non-restricted presentation of *Mycobacterium leprae* hsp 65 to $CD4^+$ T cells by transfected macrophages, 35

Mast cell function:
modulation, 508
Mucins, goblet cells:
immune-mediated alteration of terminal sugars in *Nippostrongylus brasiliensis* infection, 303
Mucosal and systemic cellular responses:
to *Toxoplasma gondii*, 421

Natural killer and LAK activity against intestinal tumour lines, 166
Natural killer cells:
and tumour necrosis factor-α-mediated enteropathy, 335
Neutrophil chemotactic desensitization and transmigration, 491
Nitrogen intermediates, reactive:
in cerebral malaria, 286

Oral tolerance:
susceptibility to, 147
Orally administered proteins:
T-cell response to, 534

Perforin and granzymes:
in cytotoxic lymphocytes, 105
Platelets:
tumoricidal activity, 650
Prostanoid:
generation by human eosinophils, 279
Protein malnutrition:
influence on secretory immune responses during development, adulthood and ageing, 308

Reshaped human antibody:
fine specificity, 364
Respiratory burst potential:
modulation by *Entamoeba histolytica* proteins, 291
Ricin:
enhancement of IgE responses, 226

Suppressive lymphokine, human $CD8^+$ $CD57^+$:
purification and partial sequence of, 205
Synovial fibroblasts:
interaction with complement C5b–9, 329

Tachykinins:
activation of human neutrophils, 629
T-cell activation:
anti-major histocompatibility complex class I-induced activation and clonal abortion, 444
T-cell activation, incomplete:
in anterior chamber-associated immune deviation, 266
T-cell receptor δ-chain transcripts:
diversity, 252
T-cell receptor expression, in Biozzi AB/H ($H\text{-}2^{dq1}$) mice, 260
T-cell responses to house dust mite allergen:
major and cryptic epitopes, 65
Thymus atrophy:
recovery from, 616
Tolerance, in Mls^a mice:
broken by anti-T-cell receptor Vβ6, 122
Toxoplasmosis: in SCID and immunocompetent mice, 430
Transforming growth factor-β_1:
effects on polymorphonuclear leucocytes, 197
Tumour necrosis factor-α:
effects of FcεRII expression, 476
induced by Theiler's murine encephalitis virus, 399
induction by extracellular matrix protein, 50

Uveoretinitis, autoimmune:
in vivo lymphokine production in, 387

Vβ usage:
in non-obese diabetic and I–E transgenic non-obese diabetic mice, 606

Xanthines:
suppression of tumour necrosis factor-α and inhibition of cAMP phosphodiesterase, 520

Zinc-binding peptides:
in *Leishmania*, 582